

## SPLIT RING SPECTRA AND SECOND PERIODICITY FAMILIES IN STABLE HOMOTOPY OF SPHERES

LIN JINKUN

(Received in revised form 9 February 1989)

### §1. INTRODUCTION

IN RECENT years, many infinite families in the stable homotopy groups of spheres  $\pi_*(S^0)_p$  had been found aside from the im  $J$ . In [15, 16], Smith proved that for any prime  $p \geq 5$ , there are  $\beta_t (t \geq 1)$  and  $\beta_{tp/j} (t \geq 1, 1 \leq j \leq p-1)$  families in  $\pi_*(S^0)_p$ , these are obtained by construction of some appropriate self maps of 4 cells spectrum  $V(1)$ . Almost at the same time, S. Oka [9] and Zahler [19] also obtained  $\beta_{tp/j} \in \pi_*(S^0)_p (t \geq 1, 1 \leq j \leq p-1)$  and [19] showed that these infinite families are detected by  $E_2$  term  $\text{Ext}_{BP_*BP}^{2,*}(BP_*, BP_*)$  of the Adams–Novikov spectral sequence based on  $BP$  theory.

$BP$  method seems to be more powerful. Miller, Ravenel and Wilson [7] gave a complete description of the generators of the above  $\text{Ext}^2$ . Their description of the generators  $\beta_{tj,j,i} (\beta_{tj,j,i} = \beta_{tj,j}$  if  $i = 1)$  suggest that if there exists certain 4 cells spectrum  $V_{j,i}$  such that  $BP_*(V_{j,i}) = BP_*/(p^i, v_1^i)$  and a self map  $f$  of  $V_{j,i}$  such that the induced  $BP_*$  homology homomorphism  $f_* = v_2^i$ , then the elements in  $\pi_*(S^0)_p$  which are detected by  $\beta_{tj,j,i}$  are obtained directly, where  $v_i$  are the generators in the coefficient ring  $BP_*$ . Since [7] reveals some periodicity phenomenas, the elements in  $\pi_*(S^0)_p$  detected by  $\beta_{tj,j,i}$  are called second periodicity elements in the literature.

Based on Toda's result on  $d_{2p-1}(\beta_{p/p}) \neq 0$ , Ravenel [13] proved that  $d_{2p-1}(\beta_{p^n/p^n}) \neq 0$  for  $n \geq 1$ , where  $d_r$  are differential in the Adams–Novikov spectral sequence. That is to say,  $\beta_{p^n/p^n}$  dies in the spectral sequence and other  $\beta_{tp^n/j} \in \text{Ext}^2$ , i.e. for  $t \neq 1$  or  $1 \leq j < p^n$ , may detect elements in  $\pi_*(S^0)_p$ . In this area, Oka did a great deal of works. Oka [9] obtained  $\beta_{tp/j} \in \pi_*(S^0)_p$  for  $p \geq 5, t \geq 1, 1 \leq j \leq p-1; j \leq p$  if  $t \geq 2$ . [10] also obtained  $\beta_{tp^2/j} \in \pi_*(S^0)_p$  for  $t \geq 1, 1 \leq j \leq 2p-2; j \leq 2p$  if  $t \geq 2$ . [11] [12] further generalized the above results to  $\beta_{tp^n/j} \in \pi_*(S^0)_p$  for  $n \geq 2, t \geq 1, 1 \leq j \leq 2^{n-1}(p-1); j \leq 2^{n-1}p$  if  $t \geq 2$ .

These are obtained on the basis of deep research on the structure of graded ring  $[\Sigma^* V_j; V_j]$ , where  $V_j$  is a 4 cells spectrum such that  $BP_*(V_j) = BP_*/(p, v_1^j)$ . [11] [12] showed that  $V_j$  is a split commutative ring spectrum if  $j \equiv 0 \pmod{p}$  and is a commutative ring spectrum but nonsplit if  $j \not\equiv 0 \pmod{p}$ . In both cases, the structure of  $[\Sigma^* V_j; V_j]$  differs in the two cases. [11] obtained the above  $\beta_{tp^n/j}$  for  $t \geq 2$  by using the structure in split case and [12] obtained the above  $\beta_{tp^n/j}$  for  $t \geq 1$  by using the structure in nonsplit case.

The main purpose of the present paper is to consider some more properties of the graded ring  $[\Sigma^* V_j; V_j]$  in split case and then we find that the above results obtained in [11] in case  $t \geq 2$  can be generalized to get almost all  $\beta_{tp^n/j} \in \pi_*(S^0)_p$ , our main results are

**THEOREM I.** For  $p \geq 5, n \geq 1, t \geq 2, 1 \leq j \leq p^n$ , the generators

$$\beta_{tp^n/j} \in \text{Ext}_{BP_*BP}^{2,*}(BP_*, BP_*)$$

survive nontrivially to  $\pi_*(S^0)_p$  in the Adams–Novikov spectral sequence and the corresponding homotopy classes are of  $p$  order and are linearly independent.

THEOREM II. If  $1 \leq j \leq p^{n-1}$  and  $j \equiv 0 \pmod{p}$ , the above  $\beta_{i p^n/j} \in \pi_*(S^0)_p$  are divisible by  $p$ .

THEOREM III. If  $p, n, t, j$  are as those in the Theorem I, then there exists a spectrum  $X$  such that  $BP_*(X) = BP_*/(p, v_1^j, v_2^{t p^n})$ .

Theorem I is proved by double induction. The input of the first induction is a result of Oka [9] that there exists a self map  $\xi_n: \Sigma^{2tp^n}(p^2-1) V_{p^n} \rightarrow V_{p^n}$  such that  $(\xi_n)_* = v_2^{t p^n}$  if  $n = 1$ . It follows from a result in [10] that  $h_1 \xi_n^p = \xi_n^p h_1$ , where  $h_{k-1}$  is a map in the following cofibration (write  $q = 2(p-1)$ )

$$\Sigma^{(k-1)p^n q} V_{p^n} \rightarrow V_{k p^n} \rightarrow V_{(k-1)p^n} \xrightarrow{h_{k-1}} \Sigma^{(k-1)p^n q+1} V_{p^n}$$

Write  $\xi_n^p = f_1$ , then  $f_1 \in [\Sigma^{t p^{n+1}(p+1)q} V_{p^n}; V_{p^n}]$  such that  $(f_1)_* = v_2^{t p^{n+1}}$ . It follows from the property of cofibration that there exists  $f_2 \in [\Sigma^{t p^{n+1}(p+1)q} V_{2 p^n}; V_{2 p^n}]$  such that  $(f_2)_* = v_2^{t p^{n+1}}$  and then start the second induction to get  $f_k \in [\Sigma^{t p^{n+1}(p+1)q} V_{k p^n}; V_{k p^n}]$  such that  $(f_k)_* = v_2^{t p^{n+1}}$  for  $1 \leq k < p$ . The second induction is done by using  $BP$  method and the properties of split ring spectra  $V_{k p^n}$  and it can be done only if  $k < p$ . By using some results on invariant ideals in  $BP_*$ , we can then prove the existence of  $\xi_{n+1}$  to complete the first induction.

## §2. SOME PRELIMINARIES ON $BP$

Let  $BP$  denote Brown–Peterson spectrum  $p$  localized,  $p$  a prime. It is well known that  $BP_* = \pi_*(BP) = Z_{(p)}[v_1, v_2, \dots]$  and  $BP_* BP = BP_*[t_1, t_2, \dots]$ , where the generators  $v_i, t_i$  have degree  $2(p^i-1)$ . An ideal  $I \subset BP_*$  is called invariant if  $I \cdot BP_* BP = BP_* BP \cdot I$ ,  $a \in BP_*$  is called mod  $I$  invariant if  $\eta_R a = \eta_L a \pmod{I \cdot BP_* BP}$ , where  $\eta_R, \eta_L: BP_* \rightarrow BP_* BP$  are right and left unit respectively. Landweber [4] proved that the only prime invariant ideal is  $I_n = (p, v_1, \dots, v_{n-1})$  ( $1 \leq n \leq \infty$ ).

For  $x \in BP_*$ , invariant ideal  $I \subset J$  of  $BP_*$ , the multiplication by  $x$  induces a  $BP_*$  homomorphism  $BP_*/I \rightarrow BP_*/J$ ,  $y \bmod I \mapsto xy \bmod J$ , which is also denoted by  $x$ . If  $x = 1$ , it is called canonical projection which is denoted by  $\rho_*$ . From [4],  $BP_*/I_n$  is a  $BP_* BP$  comodule and we have

$$\text{Ext}_{BP_* BP}^{0,*}(BP_*, BP_*/I_n) = Z_p[v_n]$$

Miller and Wilson [6] showed that  $(I_n, v_n^j, y)$  is an invariant ideal if and only if  $\bar{y} \in \text{Ext}_{BP_* BP}^{0,*}(BP_*, BP_*/(I_n, v_n^j))$ , where  $\bar{y} = y \bmod (I_n, v_n^j)$ . In the following, we give the description of the above  $\bar{y}$  given in [6] only in the  $n = 1$  case.

PROPOSITION 2.1. ([6] p. 140 Proposition 6.3 in case  $n = 1$ ) Let  $p > 2$ , then  $\text{Ext}_{BP_* BP}^{0,*}(BP_*, BP_*/(p, v_1^j))$  is a  $Z_p[v_1]/(v_1^j)$  submodule of  $BP_*/(p, v_1^j)$  generated by 1 and  $v_1^m \bar{c}_1(r)$ , where  $m = \max\{0, j - q_1(r)\}$   $r = ap^s, a \not\equiv 0 \pmod{p}$  and (cf. p. 137)

$$q_1(ap^s) = \begin{cases} p^s & \text{if } a = 1 \\ p^s + p^{s-1} - 1 & \text{otherwise} \end{cases}$$

$$\bar{c}_1(ap^s) = v_2^{ap^s} \quad \text{if } a = 1 \text{ or } s = 0$$

and for otherwise

$$\bar{c}_1(ap^s) = v_2^{ap^s} - av_1^{b_1} v_2^{a_1 p^{s-2}} - 2a \sum_{j=2}^{s-1} v_1^{b_j} v_2^{a_j p^{s-1-j}}$$

modulo  $(p, v_1^{p^s})$ , where  $a_j = (ap-1)p^j + 1$ ,  $b_j = p^s + p^{s-1} - p^{s-j} - p^{s-j-1}$  ( $j \geq 1$ ).

The following corollary is well known; the proof is omitted.

COROLLARY 2.2. If  $j \leq p^s$ ,  $v_2^{ap^s} \in \text{Ext}_{BP_*}^{0,*}(BP_*, BP_*/(p, v_1^j))$ .

The following corollary will play an important role in the proof of the main Theorem I.

COROLLARY 2.3. Let  $p > 2$ ,  $x \in BP_*$  such that  $|v_1^{(k-1)p^n} x| = tp^{n+1}(p+1)q$ , where  $q = 2(p-1)$  and  $t \geq 1$ ,  $n \geq 0$ , then

- (1) If  $2 \leq k < p$ ,  $v_1^{(k-1)p^n} x = 0$  in  $\text{Ext}_{BP_*}^{0,*}(BP_*, BP_*/(p, v_1^{kp^n}))$ .
- (2) If  $k = p$ ,  $v_1^{(k-1)p^n} x = \sum_{j=1}^{[n+1/2]} \lambda_j v_1^{p^{n+1}-p^{n+1-2j}} \bar{c}_1(a_j p^{n+1-2j})$  in  $\text{Ext}_{BP_*}^{0,*}(BP_*, BP_*/(p, v_1^{p^{n+1}}))$ , where  $\lambda_j \in \mathbb{Z}_p$ ,  $a_j = tp^{2j} - \frac{p^{2j-1}}{p-1}$ .

*Proof.* (1) Proposition 2.1 shows that  $v_1^{(k-1)p^n} x$  must be a linear combination of elements  $v_1^r \bar{c}_1(ap^s)$ , where  $(k-1)p^n \leq r < kp^n$  and  $r \geq m = \{0, kp^n - q_1(ap^s)\} \geq kp^n - p^s - p^{s-1} + 1$ . For degree reason we have

$$rq + ap^s(p+1)q = tp^{n+1}(p+1)q$$

Hence  $r = (tp^{n+1} - ap^s)(p+1)$  and so  $r$  is divisible by  $p^s$ , we have  $p$ -adic expansion of  $r$  as follows

$$r = (k-1)p^n + b_{n-1}p^{n-1} + \dots + b_s p^s$$

where  $0 \leq b_i < p$ . On the other hand  $r$  is also divisible by  $p+1$ , then

$$(k-1) - b_{n-1} + b_{n-2} - \dots + (-1)^{n-s} b_s = 0$$

and so  $b_i$  can't be all equal to  $p-1$ , i.e. there is  $j \geq s$  such that

$$b_{n-1}p^{n-1} + \dots + b_s p^s + p^s + p^{s-1} - 1 \leq (p-1)(p^{n-1} + \dots + p^s) - p^j + p^s + p^{s-1} - 1 = p^n - p^s - p^j + p^s + p^{s-1} - 1 < p^n$$

then  $r < kp^n - p^s - p^{s-1} + 1 \leq m$ , yields contradiction.

(2) If  $k = p$ , it follows from  $r = (p-1)p^n + b_{n-1}p^{n-1} + \dots + b_s p^s \geq m \geq p^{n+1} - p^s - p^{s-1} + 1$  that  $b_{n-1}p^{n-1} + \dots + b_s p^s + p^s + p^{s-1} - 1 \geq p^n$  and so  $b_{n-1}, b_{n-2}, \dots, b_s$  are all equal to  $p-1$ , and it follows from the following equation

$$(p-1) - b_{n-1} + b_{n-2} - \dots + (-1)^{n-s} b_s = 0$$

that  $n-s$  must be odd. Let  $n-s = 2j-1$  ( $j \geq 1$ ), then  $s = n+1-2j$  and  $j \leq [n+1/2]$ ,  $r = (p-1)p^n + (p-1)p^{n-1} + \dots + (p-1)p^{n+1-2j} = p^{n+1} - p^{n+1-2j}$  and from  $r = p^{n+1} - p^{n+1-2j} = (tp^{n+1} - a_j p^{n+1-2j})(p+1)$ , we have  $a_j = tp^{2j} - (p^{2j}-1)/(p-1)$ , the proposition is proved.

COROLLARY 2.4. If  $2 \leq k < p$ , then

$$\text{Ext}_{BP_*}^{0, tp^{n+1}(p+1)q - (k-1)p^n q}(BP_*, BP_*/(p, v_1^{p^n})) = 0$$

*Proof.* If there is a generator  $v_1^b \bar{c}_1(ap^s)$ , from 2.1 we have

$$p^n > b \geq m = \max \{0, p^n - q_1(ap^s)\} \geq p^n - p^s - p^{s-1} + 1$$

and for degree reason,  $bq + ap^s(p+1)q = tp^{n+1}(p+1)q - (k-1)p^nq$ . Let  $r = (k-1)p^n + b$ , then  $rq + ap^s(p+1)q = tp^{n+1}(p+1)q$  and  $(k-1)p^n \leq r < kp^n$ ,  $r \geq kp^n - p^s - p^{s-1} + 1$ . Similar to the proof of 2.3, there exists no such  $b$  in the case  $2 \leq k < p$ .

For  $t \geq 1$ , [7] defined  $\beta_{tp^n/j} = \delta \delta_j v_2^{tp^n} \in \text{Ext}_{BP_*}^{2, tp^n}(BP_*, BP_*)$ , where  $\delta, \delta_j$  are the boundary homomorphisms in the Ext sequences induced by the following two short exact sequences respectively

$$E: 0 \rightarrow BP_* \xrightarrow{p} BP_* \rightarrow BP_*/(p) \rightarrow 0$$

$$E_j: 0 \rightarrow BP_*/(p) \xrightarrow{v_1^j} BP_*/(p) \rightarrow BP_*/(p, v_1^j) \rightarrow 0$$

[7] proved that if  $1 \leq j \leq p^n$ ,  $\beta_{tp^n/j} \neq 0$ , are of  $p$  order and linearly independent, having internal degree  $2tp^n(p^2-1) - 2j(p-1)$ .

The following proposition guarantees that if there exists a self map  $f: \Sigma^{tp^n(p+1)q} V_j \rightarrow V_j$  such that  $f_* = v_1^{tp^n}$ , then  $\beta_{tp^n/j}$  survives nontrivially to  $\pi_*(S^0)_p$  in the Adams-Novikov spectral sequence, where  $V_j$  is a spectrum such that  $BP_* V_j = BP_*/(p, v_1^j)$ .

**PROPOSITION 2.5.** ([3] Theorem 1.7 or [7] Lemma 2.10). *Let  $W \rightarrow X \rightarrow Y \xrightarrow{h} \Sigma W$  be a cofibration of finite spectra such that the induced  $BP_*$  homomorphism  $h_* = 0$ ,  $\delta: \text{Ext}^s(BP_*, BP_* Y) \rightarrow \text{Ext}^{s+1}(BP_*, BP_* W)$  be boundary homomorphism in Ext exact sequence. If  $\bar{x} \in \text{Ext}^s(BP_*, BP_* Y)$  survives to  $x \in \pi_*(Y)_p$  in the Adams-Novikov spectral sequence, then  $\delta \bar{x}$  survives to  $h_*(x) \in \pi_*(W)_p$ .*

### §3. SPLIT RING SPECTRA AND ITS PROPERTIES

In this section, we recall some results on the structure of graded ring  $[\Sigma^* K; K]$  of split ring spectra  $K$  over  $Z$ , given in [11]. Moreover, we consider further properties of split ring spectra  $V_j$  over  $Z_p$  which are not in [11].

Let  $M$ , (or briefly  $M$ ) =  $S^0 \cup_i e^1$  be Moore spectrum over the cyclic group  $Z_i$ , then there is a cofibration

$$S^0 \xrightarrow{i} S^0 \xrightarrow{j_0} M \xrightarrow{\tau} \Sigma S^0$$

Since  $t \cdot 1_M$  is zero in  $[M; M]$ , then  $M \wedge M = M \vee \Sigma M$ . Let  $\mu_M: M \wedge M \rightarrow M$  be the projection to the first summand and  $\bar{\mu}_M: \Sigma M \rightarrow M \wedge M$  be injection, then define

$$d_M: [\Sigma^r M; M] \rightarrow [\Sigma^{r+1} M; M]$$

to be  $d_M(f) = \mu_M(1_M \wedge f) \bar{\mu}_M$ , proposition 1.1 in [11] showed that  $d_M$  is a derivation, i.e.  $d_M^2 = 0$  and

$$d_M(gf) = (-1)^k d_M(g)f + g d_M(f)$$

where  $g \in [\Sigma^* M; M]$ ,  $f \in [\Sigma^k M; M]$ . Let  $\delta_M = j_0 \tau: M \rightarrow \Sigma M$  then  $d_M(\delta_M) = -1_M$  (cf. [11] p. 272) and

$$[\Sigma^* M; M] = (\mu[\Sigma^* M; M] \oplus \mu[\Sigma^* M; M] \cdot \delta_M$$

where  $\mu[\Sigma^* M; M] = \ker d_M \cap [\Sigma^* M; M]$ . [11] showed that  $M$  is a commutative ring spectrum with unique multiplication  $\mu_M: M \wedge M \rightarrow M$ .

the cofibre of map  $\phi: \Sigma^k M \rightarrow M$ , i.e. there is a cofibration

$$\Sigma^k M \xrightarrow{\phi} M \xrightarrow{i} K \xrightarrow{\pi} \Sigma^{k+1} M$$

cell decomposition  $K = S^0 \cup e^1 \cup e^{k+1} \cup e^{k+2}$ , to have a decomposition  $\vee \Sigma K \vee \Sigma^{k+1} K \vee \Sigma^{k+2} K$ , we introduce

3.1. ([11] Definition 2.1)  $K$  is said to be a split ring spectrum (over  $Z_t$ ) if  $[\Sigma^* K; K]$  and  $\phi \wedge 1_K = 0$  in  $[\Sigma^k M \wedge K; M \wedge K]$ .

condition provides us with decomposition  $M \wedge K = K \vee \Sigma K$  and the second provides  $K \wedge K = (M \wedge K) \vee \Sigma^{k+1}(M \wedge K)$ , then  $K \wedge K$  splits into  $\Sigma^{k+1} K \vee \Sigma^{k+2} K$ . Let

$$\begin{aligned} \mu_1: K \wedge K &\rightarrow K, & \nu_1: K &\rightarrow K \wedge K \\ \mu_2: K \wedge K &\rightarrow \Sigma K, & \nu_2: \Sigma K &\rightarrow K \wedge K \\ \mu_3: K \wedge K &\rightarrow \Sigma^{k+1} K, & \nu_3: \Sigma^{k+1} K &\rightarrow K \wedge K \\ \mu_4: K \wedge K &\rightarrow \Sigma^{k+2} K, & \nu_4: \Sigma^{k+2} K &\rightarrow K \wedge K \end{aligned} \quad (3.2)$$

ns to each summand and injections from each summand, then  $\mu_i \nu_i = 1_K$ ,  $j$ ). Let  $d_K, d'_K$  (or briefly  $d, d'$ )

$$\begin{aligned} d_K: [\Sigma^r K; K] &\rightarrow [\Sigma^{r+1} K; K] \\ d'_K: [\Sigma^r K; K] &\rightarrow [\Sigma^{r+k+1} K; K] \end{aligned} \quad (3.3)$$

o be  $d_K(f) = \mu_1(1_K \wedge f) \nu_2, d'_K(f) = \mu_1(1_K \wedge f) \nu_3$ , then  $d_K, d'_K$  are two "derivations" on  $[\Sigma^* K; K]$ , i.e. they satisfy

$$\begin{aligned} f(g) &= (-1)^j d_K(f)g + f d_K(g) \\ f(g) &= (-1)^j d'_K(f)g + f d'_K(g) + d_K(f)(\phi' \wedge 1_K) d_K(g) \\ d_K^2 &= 0, d_K d'_K = -d'_K d_K \quad \text{and} \quad (d'_K)^2 = 0 \text{ if } t \not\equiv 0 \pmod{3} \end{aligned} \quad (3.4)$$

$\Sigma^* K; K]$  and  $\deg g = j$ ,  $\phi'$  is an element of  $\pi_*(S^0)_p$  such that  $t\phi'$  is an attaching map. ([11] p 269).

er, there exist two elements  $\delta_K \in [\Sigma^{-1} K; K]$ ,  $\delta'_K \in [\Sigma^{-k-1} K; K]$  (or briefly  $\delta, \delta'$ )

$$\begin{aligned} d(\delta) &= -1_K, & d'(\delta) &= 0, & d(\delta') &= 0, & d'(\delta') &= -1_K \\ (\delta')^2 &= 0, & \delta \delta' &= -\delta' \delta, & \delta^2 &= \delta'(\phi' \wedge 1) \\ \delta_i &= i\delta_M, & \pi \delta &= -\delta_M \pi \end{aligned} \quad (3.5)$$

$-1_K$  only holds in case  $t \not\equiv \pm 3 \pmod{9}$  (cf. [11] p. 270 and Lemma 2.6) and

$$\delta' = i\pi: K \xrightarrow{\pi} \Sigma^{k+1} M \xrightarrow{i} \Sigma^{k+1} K \quad (3.6)$$

78 (2.3)).

"derivations"  $d_K, d'_K$  make the graded ring  $[\Sigma^* K; K]$  to have the following structure. Let  $\mathcal{C}_*(K) = \ker d \cap \ker d'$  be a subgroup of  $[\Sigma^* K; K]$ , then we have

3.7. ([11] Theorem 5.5) If  $t \not\equiv 0 \pmod{3}$ ,  $\mathcal{C}_*$  is a summand of  $[\Sigma^* K; K]$  which is the direct sum in the following 16 ways

$$[\Sigma^* K; K] = \mathcal{C}_* \oplus \mathcal{D}_* \oplus \mathcal{D}'_* \oplus \mathcal{D}''_*$$

for

$$\begin{aligned}\mathcal{D}_* &= \mathcal{C}_* \delta, \delta \mathcal{C}_* \\ \mathcal{D}'_* &= \mathcal{C}_* \delta', \delta' \mathcal{C}_* \\ \mathcal{D}''_* &= \mathcal{C}_* \delta \delta', \mathcal{C}_* \delta' \delta, \delta' \mathcal{C}_* \delta, \delta \delta' \mathcal{C}_*\end{aligned}$$

Moreover, [11] Theorem 5.6 and Corollary 5.7 showed that  $\mathcal{C}_*$  is a commutative subring of  $[\Sigma^* K; K]$  and if  $t \not\equiv 0 \pmod{3}$ ,  $f \in \mathcal{C}_*$  having even degree, then  $[\delta', f'] = [\delta, f'] = 0$ , where  $[\ , \ ]$  denotes the graded commutator, i.e.  $[f, f'] = ff' - (-1)^{|f| \cdot |f'|} f'f$ , then we have

$$\delta' f' = f' \delta' \quad (3.8)$$

$$\delta f' = f' \delta \quad \text{if } f \text{ has even degree, } f \in \mathcal{C}_* \quad (3.8)$$

THEOREM 3.10. *If  $t \not\equiv 0 \pmod{3}$ , split ring spectrum  $K$  is an associative commutative ring spectrum with multiplication  $\mu_1$ . (cf. [11] Theorem 4.13)*

The following is a sufficient condition of the cofibre  $K$  of  $\phi$  to be a split ring spectrum.

PROPOSITION 3.11. ([11] Proposition 2.9). *If  $\phi = f^n (n \geq 2)$  and  $d_M(f) = 0$ , where  $f \in [\Sigma^* M; M]$  having even degree, then the cofibre  $K$  of  $\phi$  is a ring spectrum. Moreover,  $K$  is a split ring spectrum if  $n \equiv 0 \pmod{t}$ .*

The following are examples of split ring spectra given in [11], and will also be used in the proof of the main Theorems I and II.

Let  $M(m) = M_{p^m} = S^0 \cup_{p^m} e^1$  be Moore spectrum over the cyclic group  $Z_{p^m}$ , we have

THEOREM 3.12. ([11] Theorem 6.2) *There exists a self map*

$$A_m: \Sigma^{2p^{m-1}(p-1)} M(m) \rightarrow M(m), \quad m \geq 1$$

such that  $d_M(A_m) = 0$  and the induced  $BP_*$  homomorphism

$$(A_m)_* = v_1^{p^{m-1}}: BP_*/(p^m) = BP_*/(p^m)$$

Let  $M(m, sp^{m-1})$  be the cofibre of  $A_m^*$ , i.e. there is a cofibration

$$\Sigma^{sp^{m-1}q} M(m) \xrightarrow{A_m^*} M(m) \rightarrow M(m, sp^{m-1})$$

realized the following short exact sequence

$$0 \rightarrow BP_*/(p^m) \xrightarrow{v_1^{p^{m-1}}} BP_*/(p^m) \rightarrow BP_*/(p^m, v_1^{sp^{m-1}}) \rightarrow 0$$

i.e.  $BP_* M(m, sp^{m-1}) = BP_*/(p^m, v_1^{sp^{m-1}})$ , then the following corollary follows from Proposition 3.11 and Theorem 3.12 directly.

COROLLARY 3.13. *If  $s \equiv 0 \pmod{p^m}$ ,  $M(m, sp^{m-1})$  is a split ring spectrum over  $Z_{p^m}$ .*

In this paper, we only use  $M(m, sp^{m-1})$  for  $m = 1$ ,  $s = kp^n$  ( $n \geq 1$ ). Write  $V(0) = M(1) = S^0 \cup_p e^1$  be Moore spectrum over  $Z_p$ ,  $\phi = A_1: \Sigma^q V(0) \rightarrow V(0)$ ,  $V_{kp^n} = M(1, kp^n)$ , then there is a cofibration

$$\Sigma^{kp^n q} V(0) \xrightarrow{\phi^{kp^n}} V(0) \xrightarrow{i_k} V_{kp^n} \xrightarrow{\pi_k} \Sigma^{kp^n q+1} V(0) \quad (3.14)$$

realized the following short exact sequence

$$0 \rightarrow BP_*/(p) \xrightarrow{v_1^{k p^n}} BP_*/(p) \rightarrow BP_*/(p, v_1^{k p^n}) \rightarrow 0$$

Then it follows from 3.13 that if  $n \geq 1$ ,  $V_{k p^n}$  is a split ring spectrum over  $Z_p$ .

Write  $\delta'$  in (3.6) as  $\bar{h}_k$ , i.e.

$$\bar{h}_k = i_k \pi_k: V_{k p^n} \xrightarrow{\pi_k} \Sigma^{k p^n q + 1} V(0) \xrightarrow{i_k} \Sigma^{k p^n q + 1} V_{k p^n} \quad (3.15)$$

and  $\delta$  in (3.5) still write as  $\delta$ , then  $[\Sigma^* V_{k p^n}; V_{k p^n}]$  has two "derivations"  $d$  and  $d'$  and from (3.5), (3.4) we have

$$\begin{aligned} d(\bar{h}_k) &= 0, d'(\bar{h}_k) = -1_{V_{k p^n}}, d(\delta) = -1_{V_{k p^n}}, d'(\delta) = 0 \\ \delta i_k &= i_k \delta_{V(0)}, \pi_k \delta = -\delta_{V(0)} \pi_k \\ d(fg) &= (-1)^j d(f)g + fd(g) \\ d'(\bar{h}_k g) &= (-1)^j d'(\bar{h}_k)g + \bar{h}_k d'(g) \\ d'(f \bar{h}_k) &= (-1)^t d'(f) \bar{h}_k + f d'(\bar{h}_k) \end{aligned} \quad (3.16)$$

where  $j = \deg g$  and write  $\deg \bar{h}_k$  as  $t$  which is  $= -k p^n q - 1$ .

Since in this paper we only consider  $p$  as a prime  $\geq 5$ , then  $p \not\equiv 0 \pmod{3}$  and so, from (3.4) we have

$$(d')^2 = 0, d^2 = 0, dd' = -d'd \quad (3.17)$$

Let  $\mathcal{C}_*(V_{k p^n}) = \ker d \cap \ker d'$ , from Theorem 3.7 we have

$$\begin{aligned} [\Sigma^* V_{k p^n}; V_{k p^n}] &= \mathcal{C}_* \oplus \delta \mathcal{C}_* \oplus \bar{h}_k \mathcal{C}_* \oplus \bar{h}_k \delta \mathcal{C}_* \\ &= \mathcal{C}_* \oplus \mathcal{C}_* \delta \oplus \mathcal{C}_* \bar{h}_k \oplus \mathcal{C}_* \delta \bar{h}_k \end{aligned} \quad (3.18)$$

and  $\mathcal{C}_*(V_{k p^n})$  is a commutative subring. For  $f \in \mathcal{C}_*$  having even degree we have

$$\bar{h}_k f^p = f^p \bar{h}_k, \quad \delta f^p = f^p \delta \quad (3.19)$$

Now the recollection of result on split ring spectra given in [11] is concluded and we turn to consider some properties of the split ring spectrum  $V_{k p^n}$  over  $Z_p$  which are not in [11] and are important in the proof of the main theorem.

PROPOSITION 3.20. If  $r \geq k \geq 1$ ,  $m \geq n \geq 1$

$$\begin{aligned} (i_r)_*: [\Sigma^t V_{k p^n}; V(0)] &\rightarrow [\Sigma^t V_{k p^n}; V_{r p^m}] \\ (\pi_r)^*: [\Sigma^t V(0); V_{k p^n}] &\rightarrow [\Sigma^{t - r p^m q - 1} V_{r p^m}; V_{k p^n}] \end{aligned}$$

is monic, where  $i_r, \pi_r$  cf. (3.15).

In order to prove the proposition, we need the following lemma which is obvious for dimensional reason.

LEMMA (1) If  $t < -r p^m q - 2$ ,  $[\Sigma^t V_{r p^m}; V(0)] = 0$

(2) If  $t < -1$ ,  $[\Sigma^t(V(0); V_{r p^m})] = 0$ .

*Proof of Prop. 3.20.* For  $g \in [\Sigma^t V_{kp^n}; V(0)]$  such that  $i_r g = 0$ , there exists  $g_1$  such that  $g = \phi^{r p^m} g_1$  as in the following diagram

$$\begin{array}{ccccc} & & \Sigma^{r p^m q} V(0) & \xrightarrow{\phi^{r p^m}} & V(0) & \xrightarrow{i_r} & V_{r p^m} \\ & \nearrow^{g_1} & & & \nwarrow_g & & \\ & & \Sigma^t V_{kp^n} & & & & \end{array}$$

Since  $\bar{h}_k i_k g_1 = i_k \pi_k i_k g_1 = 0$ , then by acting "derivation"  $d'$  (cf. 3.16) we have

$$0 = d'(\bar{h}_k i_k g_1) = \pm d'(\bar{h}_k) i_k g_1 + \bar{h}_k d'(i_k g_1)$$

and recall that  $d'(\bar{h}_k) = -1$ , then

$$i_k(\pm g_1 + \pi_k d'(i_k g_1)) = 0$$

and so there exists  $g_2 \in [\Sigma^t V_{kp^n}; \Sigma^{r p^m q + k p^n q} V(0)]$  such that

$$\pm g_1 + \pi_k d'(i_k g_1) = \phi^{k p^n} g_2$$

Hence  $g = \phi^{r p^m} g_1 = \pm \phi^{r p^m + k p^n} g_2$  (since  $r \geq k$ ,  $m \geq n$  we have  $\phi^{r p^m} \pi_k = 0$ ). Repeating the above process, then there exists  $g_{s+1} \in [\Sigma^t V_{kp^n}; \Sigma^{r p^m q + s k p^n q} V(0)]$  such that  $g = \pm \phi^{r p^m + s k p^n} g_{s+1}$ . But it follows from Lemma 3.21 that  $g_{s+1} = 0$  for  $s$  large, so  $g = 0$ .

The proof of the second half is similar.

**PROPOSITION 3.22.** If  $k \geq 1$ ,  $n \geq 1$

$$(i_k)^*: \text{imd}' \rightarrow [\Sigma^* V(0); V_{kp^n}]$$

$$(\pi_k)_*: \text{imd}' \rightarrow [\Sigma^* V_{kp^n}; V(0)]$$

is epic.

*Proof.* For any  $g \in [\Sigma^t V(0); V_{kp^n}]$ ,  $g \pi_k \bar{h}_k = 0$  and from (3.16)

$$0 = d'(g \pi_k \bar{h}_k) = -d'(g \pi_k) \bar{h}_k + g \pi_k d'(\bar{h}_k)$$

Then  $(-d'(g \pi_k) i_k - g) \pi_k = 0$  and there exists  $g_1 \in [\Sigma^{t-k p^n q} V(0); V_{kp^n}]$  such that  $g + d'(g \pi_k) i_k = g_1 \phi^{k p^n}$ . Suppose that for  $s \geq 1$  there exists  $g_s \in [\Sigma^{t-s k p^n q} V(0); V_{kp^n}]$  such that  $g + d'(g \pi_k) i_k = g_s \phi^{s k p^n}$ , then  $0 = d'(g_s \pi_k \bar{h}_k) = -d'(g_s \pi_k) \bar{h}_k + g_s \pi_k d'(\bar{h}_k)$  and so there exists  $g_{s+1} \in [\Sigma^{t-(s+1)k p^n q} V(0); V_{kp^n}]$  such that

$$g + d'(g \pi_k) i_k = g_{s+1} \phi^{(s+1)k p^n}$$

and it follows from Lemma 3.21 that  $g_{s+1} = 0$  for  $s$  large, then  $g = -(i_k)^* d'(g \pi_k)$ . The proof of the second half is similar.

To help prove the main theorem, we now consider some cofibrations which relate  $V_{kp^n}$  and  $V_{r p^m}$ . These cofibrations can be obtained from  $3 \times 3$  Lemma (cf. [17] p.293 or [2] Lemma 6.8) in the stable homotopy category as follows.

Given a map  $\gamma = \alpha\beta: A \xrightarrow{\beta} B \xrightarrow{\alpha} C$  of spectra, there exists a cofibration

$$C(\beta) \xrightarrow{u} C(\gamma) \xrightarrow{v} C(\alpha) \xrightarrow{w} \Sigma C(\beta)$$

such that the following diagram commutes up to homotopy and sign, where  $C(x)$  denotes



the cofibre of  $\alpha$  et al

$$\begin{array}{ccccc}
 A & \xrightarrow{\gamma} & C & \xrightarrow{\quad} & C(\alpha) \xrightarrow{\quad} \Sigma C(\beta) \\
 & \searrow \beta & \nearrow \alpha & & \searrow v \\
 & & B & \xrightarrow{\quad} & C(\gamma) \xrightarrow{\quad} \Sigma B \\
 & \nearrow \beta & \searrow \alpha & & \nearrow u \\
 \Sigma^{-1} C(\alpha) & \xrightarrow{w} & C(\beta) & \xrightarrow{\quad} & \Sigma A \xrightarrow{\gamma} \Sigma C
 \end{array}$$

Recall that there is a cofibration (3.14) and let  $h_{k-1} = i_{k-1} \pi_1$ , then the following diagram commutes

$$\begin{array}{ccccccc}
 V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{p^n q+1} V_{(k-1)p^n} & \xrightarrow{\pi_{k-1}} & \Sigma^{kp^n q+2} V(0) & \xrightarrow{\phi^{kp^n}} & \Sigma^2 V(0) \\
 \downarrow \pi_1 & & \downarrow i_{k-1} & & \downarrow \psi_{k-1,k} & & \downarrow \pi_k \\
 \Sigma^{p^n q+1} V(0) & & & & C(h_{k-1}) & & \Sigma^{p^n q+2} V(0) \\
 \uparrow \phi^{(k-1)p^n} & & \uparrow \phi^{p^n} & & \uparrow i_k & & \uparrow \rho_{k,1} \\
 \Sigma^{kp^n q+1} V(0) & \xrightarrow{\phi^{kp^n}} & \Sigma V(0) & \xrightarrow{i_i} & \Sigma V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{p^n q+2} V_{(k-1)p^n}
 \end{array}$$

From the above  $3 \times 3$  lemma, we know that in the diagram  $C(h_{k-1}) = \Sigma C(\phi^{kp^n}) = \Sigma V_{kp^n}$ . Since the induced  $BP_*$  homomorphism  $(\phi^{p^n})_* = v_1^{p^n}$ , then  $(\psi_{k-1,k})_* = v_1^{p^n}$ , and the diagram commutes up to positive sign, then we have the first fundamental cofibration in this paper

$$V_{p^n} \xrightarrow{h_{k-1}} \Sigma^{p^n q+1} V_{(k-1)p^n} \xrightarrow{\psi_{k-1,k}} \Sigma V_{kp^n} \xrightarrow{\rho_{k,1}} \Sigma V_{p^n}$$

which realizes the following short exact sequence

$$0 \rightarrow BP_*/(p, v_1^{(k-1)p^n}) \xrightarrow{v_1^{p^n}} BP_*/(p, v_1^{kp^n}) \rightarrow BP_*/(p, v_1^{p^n}) \rightarrow 0$$

and also have the following relations

$$\psi_{k-1,k} i_{k-1} = i_k \phi^{p^n} \quad (3.23)$$

$$\rho_{k,1} i_k = i_1 \quad (3.24)$$

$$\pi_{k-1} = \pi_k \psi_{k-1,k} \quad (3.25)$$

$$\pi_1 \rho_{k,1} = \phi^{(k-1)p^n} \pi_k \quad (3.26)$$

Let  $h'_{k-1} = i_1 \pi_{k-1}$ , similarly we have the second fundamental cofibration as follows

$$V_{(k-1)p^n} \xrightarrow{h'_{k-1}} \Sigma^{(k-1)p^n q+1} V_{p^n} \xrightarrow{\psi_{1,k}} \Sigma V_{kp^n} \xrightarrow{\rho_{k,k-1}} \Sigma V_{(k-1)p^n}$$

which realizes a similar  $BP_*$  short exact sequence such that  $(\psi_{1,k})_* = v_1^{(k-1)p^n}$  and also have the following relations

$$\psi_{1,k} i_1 = i_k \phi^{(k-1)p^n} \quad (3.27)$$

$$\rho_{k,k-1} i_k = i_{k-1} \quad (3.28)$$

$$\pi_1 = \pi_k \psi_{1,k} \quad (3.29)$$

$$\pi_{k-1} \rho_{k,k-1} = \phi^{p^n} \pi_k \quad (3.30)$$

Let  $\gamma = \psi_{k-1,k}\psi_{1,k-1}$ , then we also have  $3 \times 3$  diagram as follows

$$\begin{array}{ccccc}
 \Sigma^{(k-1)p^n q} V_{p^n} & \xrightarrow{\gamma} & V_{kp^n} & \xrightarrow{\rho_{k,1}} & V_{p^n} \\
 \searrow \psi_{1,k-1} & & \nearrow \psi_{k-1,k} & \searrow \bar{\rho} & \nearrow v = \rho_{k-1,1} \\
 & \Sigma^{p^n q} V_{(k-1)p^n} & & C(\gamma) & \\
 \nearrow h_{k-1} & \searrow \rho_{k-1,k-2} & & \nearrow u = \psi_{k-2,k-1} & \searrow \bar{h} \\
 \Sigma^{-1} V_{p^n} & \xrightarrow{w = h_{k-2}} & \Sigma^{p^n q} V_{(k-2)p^n} & \xrightarrow{h'_{k-2}} & \Sigma^{kp^n q + 1} V_{p^n}
 \end{array}$$

It is easily seen that  $C(\gamma) = V_{(k-1)p^n}$  and  $\bar{h}\psi_{k-2,k-1} = h'_{k-2} = i_1\pi_{k-1}\psi_{k-2,k-1} = h'_{k-1}\psi_{k-2,k-1}$ . Hence we have

$$\bar{h} = h'_{k-1} + \eta\rho_{k-1,1}$$

for some  $\eta \in [\Sigma^{-(k-1)p^n q - 1} V_{p^n}; V_{p^n}]$ , and it is easily seen by degree reason that  $\eta = 0$  since  $k \geq 3$ . Therefore,  $\bar{h} = h'_{k-1}$  and so  $\gamma = \psi_{1,k}$  and  $\bar{\rho} = \rho_{k,k-1}$ . Then we have

$$\psi_{1,k} = \psi_{k-1,k}\psi_{1,k-1} = \psi_{k-1,k} \cdots \psi_{2,3}\psi_{1,2} \quad (3.31)$$

$$\rho_{k,1} = \rho_{k-1,1}\rho_{k,k-1} \quad (3.32)$$

The following relations also can be easily obtained

$$\pi_{k-j} = \pi_k \psi_{k-j,k} \quad (3.33)$$

$$\pi_j \rho_{k,j} = \phi^{(k-1)p^n} \pi_k \quad 1 \leq j < k \quad (3.34)$$

The above relations (3.23)–(3.34) will be used later.

#### §4. AN INDUCTION. PROOF OF THE MAIN THEOREM

The main theorem will be proved by double induction. The input of the first induction is the following result given in S. Oka [10].

**THEOREM 4.1.** ([10] Lemma 4.2). *If  $t \geq 2$ , there exists a self map*

$$\xi: \Sigma^{tp(p+1)q} V_p \rightarrow V_p$$

such that  $\xi_* = v_2^p: BP_*/(p, v_1^p) \rightarrow BP_*/(p, v_1^p)$  and  $\xi \in \mathcal{C}_*(V_p)$ .

For the first induction, suppose that for  $1 \leq r \leq n$  there exists

$$\xi_r: \Sigma^{tp^r(p+1)q} V_{p^r} \rightarrow V_{p^r} \text{ such that } (\xi_r)^* = v_2^{p^r} \text{ and } \xi_r \in \mathcal{C}_*(V_{p^r}).$$

Let  $f_1 = (\xi_n)^p$ , then  $(f_1)_* = v_2^{p^{n+1}}$ . In the second induction, we prove that for  $1 \leq k < p$  there exists  $f_k \in [\Sigma^{tp^{n+1}(p+1)q} V_{kp^n}; V_{kp^n}]$  such that  $(f_k)_* = v_2^{p^{n+1}}$ , and further prove the existence of  $\xi_{n+1}$  to complete the first induction.

Now we proceed to prove Theorem I. Since  $\xi_n \in \mathcal{C}_*(V_{p^n})$ , then it follows from (3.19) that  $\bar{h}_1 \xi_n^p = \xi_n^p \bar{h}_1$ , that is,  $\bar{h}_1 f_1 = f_1 \bar{h}_1$ , then there exists  $g_2$  such that the following diagram commutes

$$\begin{array}{ccccccc}
 V_{p^n} & \xrightarrow{\bar{h}_1} & \Sigma^{p^n q + 1} V_{p^n} & \xrightarrow{\psi_{1,2}} & \Sigma & V_{2p^n} & \xrightarrow{\rho_{2,1}} & \Sigma V_{p^n} \\
 \uparrow f_1 & & \uparrow f_1 & & \uparrow g_2 & & & \uparrow f_1 \\
 \Sigma^m V_{p^n} & \xrightarrow{\bar{h}_1} & \Sigma^{m+p^n q + 1} V_{p^n} & \xrightarrow{\psi_{1,2}} & \Sigma^{m+1} V_{2p^n} & \xrightarrow{\rho_{2,1}} & \Sigma^{m+1} V_{p^n}
 \end{array} \quad (4.2)$$

where  $m = tp^{n+1}(p+1)q$ . Since  $(f_1)_* = v_2^{tp^{n+1}}$ , then  $(g_2 i_0)_* = v_2^{tp^{n+1}} + v_1^{p^n} x$ , then it follows from Corollary 2.3(1) that  $(g_2)_* = v_2^{tp^{n+1}}$ . Recall that  $h_k = i_k \pi_1$ ,  $h'_k = i_1 \pi_k$ , we first prove the following

LEMMA 4.3. *There exists  $f_2 \in [\Sigma^m V_{2p^n}; V_{2p^n}]$  such that  $(f_2)_* = v_2^{tp^{n+1}}$  and  $h_2 f_1 = f_2 h_2$ ,  $h'_2 f_2 = f_1 h'_2$ .*

*Proof.* Since  $(\bar{h}_2 g_2 - g_2 \bar{h}_2) \psi_{1,2} = i_2 \pi_2 g_2 \psi_{1,2} - g_2 i_2 \pi_2 \psi_{1,2} = i_2 \pi_2 \psi_{1,2} f_1 - g_2 i_2 \pi_1 = h_2 f_1 - g_2 h_2$  and  $\rho_{2,1}(h_2 f_1 - g_2 h_2) = \rho_{2,1} i_2 \pi_1 f_1 - \rho_{2,1} g_2 i_2 \pi_1 = i_1 \pi_1 f_1 - f_1 \rho_{2,1} i_2 \pi_1 = \bar{h}_1 f_1 - f_1 \bar{h}_1 = 0$ , then there exists  $\alpha$  such that the left rectangle of the following diagram commutes

$$\begin{array}{ccccccc}
 \Sigma^{2p^n q+1} V_{p^n} & \xrightarrow{\psi_{1,2}} & \Sigma^{p^n q+1} V_{2p^n} & \xrightarrow{\rho_{2,1}} & \Sigma^{p^n q+1} V_{p^n} & \xrightarrow{\bar{h}_1} & \Sigma^{2p^n q+1} V_{p^n} \\
 \uparrow \alpha & \nearrow h_2 f_1 - g_2 h_2 & \uparrow \bar{h}_2 g_2 - g_2 \bar{h}_2 & \uparrow \beta & & & \uparrow \beta \\
 \Sigma^m V_{p^n} & \xrightarrow{\psi_{1,2}} & \Sigma^{m-p^n q} V_{2p^n} & \xrightarrow{\rho_{2,1}} & \Sigma^{m-p^n q} V_{p^n} & \xrightarrow{\bar{h}_1} & \Sigma^{m+1} V_{p^n}
 \end{array} \quad (4.4)$$

From (3.18), we can write  $\alpha = \alpha_1 + \delta \alpha_2 + \bar{h}_1 \alpha_3 + \bar{h}_1 \delta \alpha_4$  such that  $\alpha_i \in \mathcal{C}_*(V_{p^n})$  and so  $\psi_{1,2}(\alpha_1 + \delta \alpha_2) = \psi_{1,2} \alpha = h_2 f_1 - g_2 h_2$  and  $d'(\alpha_1 + \delta \alpha_2) = 0$ . Hence we may assume  $d'(\alpha) = 0$  and the above left rectangle still commutes. After we have had  $\alpha$ , then there exists  $\beta$  such that the above diagram all commutes.

Since  $\pi_1 \alpha i_1 = \pi_2 \psi_{1,2} \alpha i_1 = \pi_2 (h_2 f_1 - g_2 h_2) i_1 = 0$ , then  $\bar{h}_1 \alpha \bar{h}_1 = 0$ . Notice that  $\alpha$  has odd degree, then it follows from (3.16) that

$$0 = d'(\bar{h}_1 \alpha \bar{h}_1) = d'(\bar{h}_1) \alpha \bar{h}_1 + \bar{h}_1 d'(\alpha \bar{h}_1) = -\alpha \bar{h}_1 - \bar{h}_1 \alpha$$

then  $\bar{h}_1 \alpha = -\alpha \bar{h}_1$ .

On the other hand, it follows from the diagram (4.4) that

$$\begin{aligned}
 \pi_1 \beta \rho_{2,1} &= \pi_1 \rho_{2,1} (\bar{h}_2 g_2 - g_2 \bar{h}_2) = \phi^{p^n} \pi_2 \bar{h}_2 g_2 - \pi_1 \rho_{2,1} g_2 \bar{h}_2 = -\pi_1 f_1 i_1 \pi_2 \\
 \pi_1 \alpha \rho_{2,1} &= \pi_2 \psi_{1,2} \alpha \rho_{2,1} = \pi_2 (h_2 f_1 - g_2 h_2) \rho_{2,1} = -\pi_2 g_2 i_2 \pi_1 \rho_{2,1} \\
 &= -\pi_2 g_2 i_2 \phi^{p^n} \pi_2 = -\pi_2 g_2 \psi_{1,2} i_1 \pi_2 = -\pi_1 f_1 i_1 \pi_2
 \end{aligned}$$

Then  $\pi_1 \beta \rho_{2,1} = \pi_1 \alpha \rho_{2,1}$  and there exists  $\gamma \in [\Sigma^m V_{p^n}; \Sigma^{2p^n q+1} V(0)]$  such that  $\pi_1 \beta = \pi_1 \alpha + \gamma \bar{h}_1$ . Then  $\bar{h}_1 \beta = \bar{h}_1 \alpha + i_1 \gamma \bar{h}_1$ . From diagram (4.4) we have  $\bar{h}_1 \beta = \alpha \bar{h}_1$ , then

$$2\alpha \bar{h}_1 = i_1 \gamma \bar{h}_1$$

By acting the "derivation"  $d'$  (using (3.16)) we have  $-2\alpha = -d'(i_1 \gamma) \bar{h}_1 - i_1 \gamma$ , thus  $2\bar{h}_1 \alpha = \bar{h}_1 d'(i_1 \gamma) \bar{h}_1$ . By acting  $d'$  again, we have

$$2\alpha = d'(i_1 \gamma) \bar{h}_1 - \bar{h}_1 d'(i_1 \gamma)$$

Let

$$f_2 = g_2 + \frac{1}{2} \psi_{1,2} d'(i_1 \gamma) \rho_{2,1}$$

then

$$\begin{aligned}
 f_2 h_2 &= g_2 h_2 + \frac{1}{2} \psi_{1,2} d'(i_1 \gamma) i_1 \pi_1 \\
 &= g_2 h_2 + \psi_{1,2} \alpha \\
 &= g_2 h_2 + (h_2 f_1 - g_2 h_2) = h_2 f_1
 \end{aligned}$$

Moreover,  $(\bar{h}_2 f_2 - f_2 \bar{h}_2) \psi_{1,2} = \bar{h}_2 f_2 \psi_{1,2} - f_2 i_2 \pi_2 \psi_{1,2} = h_2 f_1 - f_2 h_2 = 0$ , then there

exists  $\eta$  such that the following diagram commutes

$$\begin{array}{ccccccc}
 \Sigma^{2p^n q+1} V_{p^n} & \xrightarrow{\psi_{1,2}} & \Sigma^{p^n q+1} V_{2p^n} & \xrightarrow{\rho_{2,1}} & \Sigma^{p^n q+1} V_{p^n} & \xrightarrow{\bar{h}_1} & \Sigma^{2p^n q+2} V_{p^n} \\
 \uparrow 0 & & \uparrow \bar{h}_2 f_2 - f_2 \bar{h}_2 & & \uparrow \eta & & \uparrow 0 \\
 \Sigma^m V_{p^n} & \xrightarrow{\psi_{1,2}} & \Sigma^{m-p^n q} V_{2p^n} & \xrightarrow{\rho_{2,1}} & \Sigma^{m-p^n q} V_{p^n} & \xrightarrow{\bar{h}_1} & \Sigma^{m+1} V_{p^n}
 \end{array}$$

By acting  $d'$  on  $\bar{h}_1 \eta = 0$  we have  $\eta = -\bar{h}_1 d'(\eta)$ . Since  $\pi_1 \eta i_1 = \pi_1 \eta \rho_{2,1} i_2 = \pi_1 \rho_{2,1} (\bar{h}_2 f_2 - f_2 \bar{h}_2) i_2 = 0$ , then  $\bar{h}_1 \eta \bar{h}_1 = 0$  and by acting  $d'$  again we have  $-\eta \bar{h}_1 - \bar{h}_1 \eta + \bar{h}_1 d'(\eta) \bar{h}_1 = 0$ , i.e.  $-2\eta \bar{h}_1 = 0$ , then we have  $\eta = -d'(\eta) \bar{h}_1$  and  $h'_2 f_2 - f_1 h'_2 = \rho_{2,1} (\bar{h}_2 f_2 - f_2 \bar{h}_2) = \eta \rho_{2,1} = 0$ .

Since the induced  $BP_*$  homomorphism  $(d'(i_1 \gamma) i_0)_* \in \text{Ext}_{BP_* BP_*}^{0, m-p^n q}(BP_*, BP_*/(p, v_1^{p^n})) = 0$  (cf. Cor. 2.4), where  $i_0: S^0 \rightarrow V_{p^n}$  is the injection of the bottom cell, then  $(d'(i_1 \gamma))_* = 0: BP_*/(p, v_1^{p^n}) \rightarrow BP_*/(p, v_1^{p^n})$  and so  $(f_2)_* = (g_2)_* = v_2^{p^n+1}$ . The lemma is proved.

Now we begin to do the second induction. As an induction hypothesis, suppose that for  $3 \leq k < p$ , there exists  $f_{k-1} \in [\Sigma^m V_{(k-1)p^n}; V_{(k-1)p^n}]$  such that  $(f_{k-1})_* = v_2^{p^n+1}$  and

$$h_{k-1} f_1 = f_{k-1} h_{k-1}, \quad h'_{k-1} f_{k-1} = f_1 h'_{k-1} \quad (4.5)$$

$$\psi_{1,k-1} f_1 = f_{k-1} \psi_{1,k-1}, \quad \rho_{k-1,1} f_{k-1} = f_1 \rho_{k-1,1} \quad (4.6)$$

If  $k = 3$ , the above condition holds from Lemma 4.3 and (4.2).

Then there exist  $g_k, g'_k$  such that the following diagrams commute

$$\begin{array}{ccccccc}
 V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{p^n q+1} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma V_{kp^n} & \xrightarrow{\rho_{k,1}} & \Sigma V_{p^n} \\
 \uparrow f_1 & & \uparrow f_{k-1} & & \uparrow g_k & & \uparrow f_1 \\
 \Sigma^m V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{m+p^n q+1} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma^{m+1} V_{kp^n} & \xrightarrow{\rho_{k,1}} & \Sigma^{m+1} V_{p^n}
 \end{array} \quad (4.7)$$

$$\begin{array}{ccccccc}
 V_{(k-1)p^n} & \xrightarrow{h'_{k-1}} & \Sigma^{(k-1)p^n q+1} V_{p^n} & \xrightarrow{\psi_{1,k}} & \Sigma V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & \Sigma V_{(k-1)p^n} \\
 \uparrow f_{k-1} & & \uparrow f_1 & & \uparrow g'_k & & \uparrow f_{k-1} \\
 \Sigma^m V_{(k-1)p^n} & \xrightarrow{h'_{k-1}} & \Sigma^{m+(k-1)p^n q+1} V_{p^n} & \xrightarrow{\psi_{1,k}} & \Sigma^{m+1} V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & \Sigma^{m+1} V_{(k-1)p^n}
 \end{array} \quad (4.8)$$

Since  $(f_{k-1})_* = v_2^{p^n+1}$ , then  $(g'_k)_* = v_2^{p^n+1} + v_1^{(k-1)p^n} x$ , it follows from Corollary 2.3(1) that  $(g'_k)_* = v_2^{p^n+1}$ .

It follows from diagram (4.7) that  $(g_k)_* = v_2^{p^n+1} + v_1^{p^n} x$  for some  $x \in BP_*$  and  $v_1^{p^n} (g_k)_* = v_1^{p^n} v_2^{p^n+1}$ , then  $v_1^{2p^n} x = 0 \pmod{(p, v_1^{k p^n})}$ , then it follows from Corollary 2.3(1) that  $(g_k)_* = v_2^{p^n+1}$ .

Now it follows from (4.7) and (4.6) that  $g_k \psi_{1,k} = g_k \psi_{k-1,k} \psi_{1,k-1} = \psi_{k-1,k} f_{k-1} \psi_{1,k-1} = \psi_{1,k} f_1$  and from (4.8), (4.6) that  $\rho_{k,1} g'_k = \rho_{k-1,k} \rho_{k,k-1} g'_k = \rho_{k-1,1} f_{k-1} \rho_{k,k-1} = f_1 \rho_{k,1}$ , then there exist  $f'_{k-1}$  and  $f''_{k-1}$  such that the following diagrams commute

$$\begin{array}{ccccccc}
 V_{(k-1)p^n} & \xrightarrow{h'_{k-1}} & \Sigma^{(k-1)p^n q+1} V_{p^n} & \xrightarrow{\psi_{1,k}} & \Sigma V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & \Sigma V_{(k-1)p^n} \\
 \uparrow f'_{k-1} & & \uparrow f_1 & & \uparrow g_k & & \uparrow f'_{k-1} \\
 \Sigma^m V_{(k-1)p^n} & \xrightarrow{h'_{k-1}} & \Sigma^{m+(k-1)p^n q+1} V_{p^n} & \xrightarrow{\psi_{1,k}} & \Sigma^{m+1} V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & \Sigma^{m+1} V_{(k-1)p^n}
 \end{array} \quad (4.9)$$

$$\begin{array}{ccccccc}
 V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{p^n q + 1} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma V_{kp^n} & \xrightarrow{\rho_{k,1}} & \Sigma V_{p^n} \\
 \uparrow f_1 & & \uparrow f'_{k-1} & & \uparrow g'_k & & \uparrow f_1 \\
 \Sigma^m V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{m+p^n q + 1} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma^{m+1} V_{kp^n} & \xrightarrow{\rho_{k,1}} & \Sigma^{m+1} V_{p^n}
 \end{array} \quad (4.10)$$

From (4.8) and (4.9), we have  $h'_{k-1}(f'_{k-1} - f_{k-1}) = 0$  and  $\rho_{k,k-1}(g_k - g'_k) = (f'_{k-1} - f_{k-1})\rho_{k,k-1}$ , then  $0 = \rho_{k-1}(g_k - g'_k) = \rho_{k-1,1}(f'_{k-1} - f_{k-1})\rho_{k,k-1}$ . Then it follows from the following Lemma 4.13 that

$$(f'_{k-1} - f_{k-1})h_{k-1} = 0 \quad (4.11)$$

From (4.7), (4.10) we have  $(f_{k-1} - f''_{k-1})h_{k-1} = 0$  and  $\psi_{k-1,k}(f_{k-1} - f''_{k-1}) = (g_k - g'_k)\psi_{k-1,k}$ , then  $\psi_{k-1,k}(f_{k-1} - f''_{k-1})\psi_{1,k-1} = (g_k - g'_k)\psi_{1,k} = 0$ , then it follows from Lemma 4.13 that

$$h'_{k-1}(f_{k-1} - f''_{k-1}) = 0. \quad (4.12)$$

(4.11) and (4.12) will be used later.

The proof of the following lemma will be postponed to the last of the proof of main theorem.

LEMMA 4.13: (1) For  $g \in [\Sigma^i V_{(k-1)p^n}; V_{(k-1)p^n}]$  such that  $\rho_{k-1,1}g i_{k-1} = 0$  and  $h'_{k-1}g = 0$ , then  $gh_{k-1} = 0$ .

(2) For  $g$  as above such that  $\psi_{k-1,k}g\psi_{1,k-1} = 0$  and  $gh_{k-1} = 0$ , then  $h'_{k-1}g = 0$ .

Now we see that  $(\bar{h}_k g_k - g'_k \bar{h}_k)\psi_{1,k} = i_k \pi_k g_k \psi_{1,k} - g'_k i_k \pi_k \psi_{1,k} = i_k \pi_k \psi_{1,k} f_1 - g'_k i_k \pi_1 = h_k f_1 - g'_k h_k$  and  $\rho_{k,k-1}(h_k f_1 - g'_k h_k) = \rho_{k,k-1} i_k \pi_1 f_1 - \rho_{k,k-1} g'_k i_k \pi_1 = i_{k-1} \pi_1 f_1 - f_{k-1} i_{k-1} \pi_1 = 0$  (cf. inductive hypothesis (4.5)). Then similar to the proof of Lemma 4.3 there exists  $\alpha$  such that  $d'(\alpha) = 0$  and there exists  $\beta$  such that the following diagram commutes

$$\begin{array}{ccccccc}
 \Sigma^{kp^n q + 1} V_{p^n} & \xrightarrow{\psi_{1,k}} & \Sigma^{p^n q + 1} V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & \Sigma^{p^n q + 1} V_{(k-1)p^n} & \xrightarrow{h'_{k-1}} & \Sigma^{kp^n q + 2} V_{p^n} \\
 \uparrow \alpha & \nearrow h_k f_1 - g'_k h_k & \uparrow \bar{h}_k g_k - g'_k \bar{h}_k & & \uparrow \beta & & \uparrow \bar{x} \\
 \Sigma^m V_{p^n} & \xrightarrow{\psi_{1,k}} & \Sigma^{m-(k-1)p^n q} V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & \Sigma^{m-(k-1)p^n q} V_{(k-1)p^n} & \longrightarrow & \Sigma^{m+1} V_{p^n}
 \end{array} \quad (4.14)$$

On the other hand,  $\rho_{k,1}(\bar{h}_k g_k - g'_k \bar{h}_k)\psi_{k-1,k} = \rho_{k,1} i_k \pi_k g_k \psi_{k-1,k} - \rho_{k,1} g'_k i_k \pi_k \psi_{k-1,k} = i_1 \pi_k \psi_{k-1,k} f_{k-1} - f_1 i_1 \pi_k \psi_{k-1,k} = h'_{k-1} f_{k-1} - f_1 h'_{k-1} = 0$ , then there exists  $\bar{\alpha}, \bar{\beta}$  such that the following diagram commutes

$$\begin{array}{ccccccc}
 \Sigma^{kp^n q + 1} V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{(k+1)p^n q + 2} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma^{kp^n q + 2} V_{kp^n} & \xrightarrow{\rho_{k,1}} & \Sigma^{kp^n q + 2} V_{p^n} \\
 \uparrow \bar{\alpha} & & \uparrow \bar{\beta} & & \uparrow \bar{h}_k g_k - g'_k \bar{h}_k & & \uparrow \bar{z} \\
 \Sigma^m V_{p^n} & \xrightarrow{h_{k-1}} & \Sigma^{m+p^n q + 1} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma^{m+1} V_{kp^n} & \xrightarrow{\rho_{k,1}} & \Sigma^{m+1} V_{p^n}
 \end{array} \quad (4.15)$$

From diagram (4.14),  $\pi_1 \alpha = \pi_k \psi_{1,k} \alpha = \pi_k (h_k f_1 - g'_k h_k) = -\pi_k g'_k i_k \pi_1 = -\pi_k g'_k i_k \pi_{k-1} \psi_{1,k-1}$ ,

then there exists  $\eta$  such that the following diagram commutes

$$\begin{array}{ccccccc}
 \Sigma^{kp^nq+1}V_{p^n} & \xrightarrow{\pi_1} & \Sigma^{(k+1)p^nq+2}V(0) & \xrightarrow{\phi^{p^n}} & \Sigma^{kp^nq+2}V(0) & \xrightarrow{i_1} & \Sigma^{kp^nq+2}V_{p^n} \\
 \uparrow -\alpha & & \uparrow \pi_k g'_k i_k \pi_{k-1} & & \uparrow \eta & & \uparrow -\alpha \\
 \Sigma^m V_{p^n} & \xrightarrow{\psi_{1,k-1}} & \Sigma^{m-(k-2)p^nq}V_{(k-1)p^n} & \xrightarrow{\rho_{k-1,k-2}} & \Sigma^{m-(k-2)p^nq}V_{(k-2)p^n} & \xrightarrow{h'_{k-2}} & \Sigma^{m+1}V_{p^n}
 \end{array} \quad (4.16)$$

Since

$$\begin{aligned}
 \phi^{p^n} \pi_k g'_k i_k \pi_{k-1} &= \pi_{k-1} \rho_{k,k-1} g'_k i_k \pi_{k-1} = \pi_{k-1} f_{k-1} \rho_{k,k-1} i_k \pi_{k-1} \\
 &= \pi_{k-1} f_{k-1} i_{k-1} \pi_{k-1} = \pi_k \psi_{k-1,k} f_{k-1} i_{k-1} \pi_{k-1} \\
 &= \pi_k g_k \psi_{k-1,k} i_{k-1} \pi_{k-1} = \pi_k g_k i_k \phi^{p^n} \pi_{k-1} = \pi_k g_k i_k \pi_{k-2} \rho_{k-1,k-2},
 \end{aligned}$$

then

$$\eta \rho_{k-1,k-2} = \pi_k g_k i_k \pi_{k-2} \rho_{k-1,k-2}$$

and there exists  $\gamma \in [\Sigma^{m+1}V_{p^n}, \Sigma^{kp^nq+2}V(0)]$  such that

$$\eta = \pi_k g_k i_k \pi_{k-2} + \gamma i_1 \pi_{k-2}$$

and  $-\alpha i_1 \pi_{k-2} = i_1 \eta = i_1 \pi_k g_k i_k \pi_{k-2} + i_1 \gamma i_1 \pi_{k-2}$ , then we have  $i_1 \pi_k g_k i_k \pi_1 = -\alpha \bar{h}_1 - i_1 \gamma \bar{h}_1$  and

$$i_1 \pi_k g_k i_k \pi_1 = -\alpha \bar{h}_1 + \bar{h}_1 d'(i_1 \gamma) \bar{h}_1 \quad (4.17)$$

since  $0 = d'(\bar{h}_1 i_1 \gamma) = i_1 \gamma + \bar{h}_1 d'(i_1 \gamma)$ .

But  $-\alpha \bar{h}_1 = \bar{h}_1 \alpha = -i_1 \pi_k g'_k i_k \pi_1$ , then  $i_1 \pi_k (g_k + g'_k) i_k \pi_1 = i_1 \pi_1 d'(i_1 \gamma) i_1 \pi_1$  and so

$$i_1 \pi_k (g_k + g'_k - \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) i_k \pi_1 = 0 \quad (4.18)$$

Recall to (4.11),  $\rho_{k,k-1} (g_k - g'_k) i_k \pi_1 = (f'_{k-1} - f_{k-1}) \rho_{k,k-1} i_k \pi_1 = (f'_{k-1} - f_{k-1}) h_{k-1} = 0$ , then there exists  $\varepsilon$  such that  $d'(\varepsilon) = 0$  and the following diagram commutes

$$\begin{array}{ccccc}
 \Sigma^{(k-1)p^nq}V_{p^n} & \xrightarrow{\psi_{1,k}} & V_{kp^n} & \xrightarrow{\rho_{k,k-1}} & V_{(k-1)p^n} \\
 & & \uparrow g_k - g'_k - \psi_{1,k} d'(i_1 \gamma) \rho_{k,1} & & \\
 & & \Sigma^m V_{kp^n} & & \\
 & & \uparrow i_k & & \\
 & & \Sigma^m V(0) & & \\
 & & \uparrow \pi_1 & & \\
 & & \Sigma^{m-p^nq-1}V_{p^n} & & \\
 \varepsilon \swarrow & & & & \searrow 0
 \end{array}$$

It is easily seen that  $\pi_1 \varepsilon i_1 = \pi_k \psi_{1,k} \varepsilon i_1 = 0$ , then  $\bar{h}_1 \varepsilon \bar{h}_1 = 0$  and then  $\bar{h}_1 \varepsilon = -\varepsilon \bar{h}_1$ . On the other hand

$$\begin{aligned}
 -\varepsilon \bar{h}_1 &= \bar{h}_1 \varepsilon = i_1 \pi_k \psi_{1,k} \varepsilon = i_1 \pi_k (g_k - g'_k - \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) i_k \pi_1 \\
 &= 2i_1 \pi_k (g_k - \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) i_k \pi_1 \quad (\text{summing with (4.18)}) \\
 &= 2i_1 \pi_k g_k i_k \pi_1 - 2\bar{h}_1 d'(i_1 \gamma) \bar{h}_1 \\
 &= -2\alpha \bar{h}_1 \quad (\text{from (4.17)})
 \end{aligned}$$

Then by acting  $d'$  we have  $\varepsilon = 2\alpha$  and

$$\begin{aligned}(g_k - g'_k - \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) h_k &= \psi_{1,k} \varepsilon = 2\psi_{1,k} \alpha \\ &= 2(h_k f_1 - g'_k h_k)\end{aligned}$$

Hence if we let

$$f_k = \frac{1}{2}(g_k + g'_k - \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) \quad (4.19)$$

we have  $f_k h_k = h_k f_1$ .

Similarly, recall to (4.12),  $i_1 \pi_k (g_k - g'_k) \psi_{k-1,k} = i_1 \pi_k \psi_{k-1,k} (f_{k-1} - f''_{k-1}) = h'_{k-1} (f_{k-1} - f''_{k-1}) = 0$ , then there exists  $\bar{\varepsilon}$  with  $d'(\bar{\varepsilon}) = 0$  such that the following diagram commutes

$$\begin{array}{ccccc} & & \Sigma^{k p^n q + 1} V_{p^n} & & \\ & \nearrow 0 & \uparrow i_1 & \nwarrow \bar{\varepsilon} & \\ & & \Sigma^{k p^n q + 1} V(0) & & \\ & & \uparrow \pi_k & & \\ & & V_{k p^n} & & \\ & & \uparrow g_k - g'_k + \psi_{1,k} d'(i_1 \gamma) \rho_{k,1} & & \\ \Sigma^{m + p^n q} V_{(k-1)p^n} & \xrightarrow{\psi_{k-1,k}} & \Sigma^m V_{k p^n} & \xrightarrow{\rho_{k,1}} & \Sigma^m V_{p^n} \end{array}$$

i.e.  $\bar{\varepsilon} \rho_{k,1} = i_1 \pi_k (g_k - g'_k + \psi_{1,k} d'(i_1 \gamma) \rho_{k,1})$  and

$$\begin{aligned}\bar{\varepsilon} \bar{h}_1 &= \bar{\varepsilon} \rho_{k,1} i_k \pi_1 = i_1 \pi_k (g_k - g'_k + \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) i_k \pi_1 \\ &= 2i_1 \pi_k g_k i_k \pi_1 \quad (\text{by summing with (4.18)})\end{aligned}$$

Since  $\bar{\alpha} \bar{h}_1 = \bar{\alpha} i_1 \pi_1 = \bar{\alpha} \rho_{k,1} i_k \pi_1 = \rho_{k,1} (\bar{h}_k g_k - g'_k \bar{h}_k) i_k \pi_1 = i_1 \pi_k g_k i_k \pi_1$ , then  $\bar{\varepsilon} \bar{h}_1 = 2\bar{\alpha} \bar{h}_1$  and by acting  $d'$  we have

$$\bar{\varepsilon} = -d'(\bar{\varepsilon} \bar{h}_1) = -2d'(\bar{\alpha} \bar{h}_1) = 2\bar{\alpha} - 2d'(\bar{\alpha}) \bar{h}_1$$

and so

$$\begin{aligned}h'_k (g_k - g'_k + \psi_{1,k} d'(i_1 \gamma) \rho_{k,1}) &= \bar{\varepsilon} \rho_{k,1} = 2\bar{\alpha} \rho_{k,1} \\ &= 2\rho_{k,1} (\bar{h}_k g_k - g'_k \bar{h}_k) = 2(h'_k g_k - f_1 h'_k)\end{aligned}$$

Hence  $f_1 h'_k = h'_k f_k$  (cf. (4.19)).

Since  $k < p$ , the induced  $BP_*$  homomorphism  $(d'(i_1 \gamma) i_0)_* \in \text{Ext}_{BP_* BP_*}^{0, m-(k-1)p^n q}(BP_*, BP_*/(p, v_1^{p^n})) = 0$  (cf. Corollary 2.4), then  $d'(i_1 \gamma)_* = 0$  and  $(f_k)_* = \frac{1}{2}[(g_k)_* + (g'_k)_*] = v_2^{p^n+1}$ . Recall to (4.19), it is easily seen that  $f_k \psi_{1,k} = \psi_{1,k} f_1$ ,  $\rho_{k,1} f_k = f_1 \rho_{k,1}$ , then the second induction is completed.

Hence we get a self map  $f_{p-1}: \Sigma^m V_{(p-1)p^n} \rightarrow V_{(p-1)p^n}$  such that  $(f_{p-1})_* = v_2^{p^n+1}$  and  $h_{p-1} f_1 = f_{p-1} h_{p-1}$ ,  $h'_{p-1} f_{p-1} = f_1 h'_{p-1}$ , then there exists  $g'_p: \Sigma^m V_{p^n+1} \rightarrow V_{p^n+1}$  such that  $\rho_{p,p-1} g'_p = f_{p-1} \rho_{p,p-1}$ . Then  $(g'_p)_* = v_2^{p^n+1} + v_1^{(p-1)p^n} x$  for some  $x \in BP_*$ . It follows from Corollary 2.3(2) that

$$v_1^{(p-1)p^n} x = \sum_{j=1}^{[n+1/2]} \lambda_j v_1^{p^n+1-p^n+1-2j} \bar{c}_1(a_j p^{n+1-2j})$$

where  $\lambda_j \in \mathbb{Z}_p$  and  $a_j \geq 2$ .

The proof of the following lemma will be postponed to the last of the proof of main theorem.

LEMMA 4.22. Suppose that for  $a \geq 2, 0 \leq j \leq s, v_2^{ap^j}$  can be realized in  $[\Sigma^* V_{p^j}; V_{p^j}]$ , then  $\bar{c}_1(ap^s)$  can be realized in  $[\Sigma^* V_{p^s}; V_{p^s}]$  and  $v_1^{p^n - p^s} \bar{c}_1(ap^s)$  can be realized in  $[\Sigma^* V_{p^n}; V_{p^n}]$ .

From the first inductive hypothesis, for  $k \leq n, v_2^{p^k}$  can be realized by  $\xi_k$  in  $[\Sigma^* V_{p^k}; V_{p^k}]$ , then it follows from Lemma 4.22 that for  $1 \leq j \leq [n + 1/2]$ ,  $v_1^{p^{n+1} - 2^j} \bar{c}_1(a_j p^{n+1-2^j})$  can be realized in  $[\Sigma^* V_{p^{n+1}}; V_{p^{n+1}}]$ , then there exists  $\bar{g} \in [\Sigma^m V_{p^{n+1}}; V_{p^{n+1}}]$  such that  $(\bar{g})_* = v_1^{(p-1)p^n} x$  and then  $(\xi_{n+1})_* = v_2^{p^{n+1}}$  if we put  $\xi_{n+1} = g'_p - \bar{g}$ . Then the first induction is also completed.

Now if  $j \leq p^n$ , let  $\rho: V_{p^n} \rightarrow V_j$  be projection, then  $\rho \xi_n i_0: \Sigma^{ip^n(p+1)q} S^0 \rightarrow V_j$  induces  $BP_*$  homomorphism  $(\rho \xi_n i_0)_* = v_2^{p^n}$ . Since  $\rho \xi_n i_0$  can be extended to  $\eta_j: \Sigma^{ip^n(p+1)q} V_j \rightarrow V_j$  such that  $(\eta_j)_* = v_2^{p^n}$ , then it follows from Proposition 2.5 that  $\beta_{ip^n/j} = \delta \delta_j v_2^{ip^n}$  survives to the following nontrivial map

$$\Sigma^{ip^n(p+1)q} S^0 \xrightarrow{i_0} \Sigma^{ip^n(p+1)q} V_j \xrightarrow{\eta_j} V_j \xrightarrow{\pi} \Sigma^{jq+1} V(0) \xrightarrow{\tau} \Sigma^{jq+2} S^0$$

the main Theorem I is then proved.

In the following, we give the proof of Lemma 4.13 and 4.22.

*Proof of Lemma 4.13.* (1) Since  $h'_{k-1}g = i_1 \pi_{k-1}g = 0$ , then  $i_1 \pi_{k-1}g i_{k-1} \pi_1 = 0$ . It follows from Proposition 3.20 that  $\pi_{k-1}g i_{k-1} \pi_1 = 0$ . Then there exists  $u \in [\Sigma^i V_{p^n}; \Sigma^{p^n q+1} V(0)]$  such that  $g i_{k-1} \pi_1 = i_{k-1} u$ , and from Proposition 3.22, there exists  $\bar{u} \in [\Sigma^i V_{p^n}; V_{p^n}]$  such that  $d'(\bar{u}) = 0$  and  $u = \pi_1 \bar{u}$  and so

$$\bar{h}_1 \bar{u} = i_1 \pi_1 \bar{u} = \rho_{k-1,1} i_{k-1} \pi_1 \bar{u} = \rho_{k-1,1} g i_{k-1} \pi_1 = 0$$

Then  $\bar{u} = d'(\bar{h}_1 \bar{u}) = 0$  and so  $g i_{k-1} \pi_1 = 0$ .

The proof of the second half is similar.

*Proof of Lemma 4.22.* From Proposition 2.1, we have

$$\bar{c}_1(ap^s) = v_2^{ap^s} - av_1^{b_1} v_2^{a_1 p^{s-2}} - 2a \sum_{j=2}^{s-2} v_1^{b_j} v_2^{a_j p^{s-1-j}} \bmod(p, v_1^{p^s})$$

where  $b_j = p^s + p^{s-1} - p^{s-j} - p^{s-1-j}$  and  $a_j = (ap-1)p^j + 1, j \geq 1$ . From the hypothesis,  $v_2^{ap^s}$  can be realized in  $[\Sigma^* V_{p^s}; V_{p^s}]$  and  $v_2^{a_j p^{s-1-j}}$  can be realized by a self map  $\varepsilon_j$  of  $V_{p^{s-1-j}}$ . Consider the cofibration

$$\Sigma^{b_j q} V_{p^{s-1-j}} \xrightarrow{\bar{\psi}} V_{p^s + p^{s-1} - p^{s-j} - p^{s-1-j}} \xrightarrow{\bar{\rho}} V_{b_j} \xrightarrow{h} \Sigma^{b_j q+1} V_{p^{s-1-j}}$$

in which  $(\bar{\psi})_* = v_1^{b_j}$  and let  $\rho': V_{p^s + p^{s-1} - p^{s-j} - p^{s-1-j}} \rightarrow V_{p^s}$ , then  $(\rho' \bar{\psi} \varepsilon_j i_0)_* = v_1^{b_j} v_2^{a_j p^{s-1-j}} \in \text{Ext}^0(BP_*, BP_*/(p, v_1^{p^s}))$ . Since  $\rho' \bar{\psi} \varepsilon_j i_0$  can be extended to a self map of  $V_{p^s}$  which has the same  $BP_*$  homomorphism, then  $v_1^{b_j} v_2^{a_j p^{s-1-j}}$  can be realized by a self map of  $V_{p^s}$ , then  $\bar{c}_1(ap^s)$  can be realized by a self map of  $V_{p^s}$ . By a similar method we can prove that  $v_1^{p^n - p^s} \bar{c}_1(ap^s)$  can be realized in  $[\Sigma^* V_{p^n}; V_{p^n}]$ . The lemma is then proved.

*Proof of Theorem II.* Recall that we have a cofibration

$$S^0 \xrightarrow{p} S^0 \xrightarrow{j_0} V(0) \xrightarrow{\tau} \Sigma S^0$$

and  $\delta_{V(0)} = j_0 \tau: V(0) \rightarrow \Sigma V(0)$  (we will write  $\delta_{V(0)}$  as  $\delta$  if there is no confusion). Let  $C(\delta)$  be the cofibre of  $\delta$ , then it follows from  $3 \times 3$  lemma that there is a commutative diagram as



follows

$$\begin{array}{ccccccc}
 \Sigma^{-1} V(0) & \xrightarrow{\delta} & V(0) & \xrightarrow{\tau} & \Sigma S^o & \xrightarrow{p^2} & \Sigma S^o \\
 & \searrow \tau & \nearrow j_0 & \searrow \lambda & \nearrow p & \searrow p & \nearrow \Sigma S^o \\
 & & S^o & & C(\delta) & & S^o \\
 & \nearrow p & \searrow p & \nearrow j' & \searrow \rho & \nearrow \tau & \searrow j_0 \\
 S^o & \xrightarrow{p^2} & S^o & \xrightarrow{j_0} & V(0) & \xrightarrow{\delta} & \Sigma V(0)
 \end{array}$$

and we find that  $c(\delta) = C(p^2)$  which we write as  $M(p^2)$ , i.e. there are cofibrations

$$S^o \xrightarrow{p^2} S^o \xrightarrow{j'} M(p^2) \xrightarrow{\tau'} \Sigma S^o$$

$$\Sigma^{-1} V(0) \xrightarrow{\delta} V(0) \xrightarrow{\lambda} M(p^2) \xrightarrow{\rho} V(0)$$

which realize the following short exact sequences

$$0 \longrightarrow BP_* \xrightarrow{p^2} BP_* \xrightarrow{j_*} BP_*/(p^2) \longrightarrow 0$$

$$0 \longrightarrow BP_*/(p) \xrightarrow{\lambda_* = p} BP_*/(p^2) \xrightarrow{\rho_*} BP_*/(p) \longrightarrow 0$$

and we have  $\tau = \tau' \lambda$ ,  $j_0 = j' \rho$ .

From (3.5) we have  $\delta_{V_j}: V_j \rightarrow \Sigma V_j$  such that  $\delta_{V_j} i = i \delta_{V(0)}$ , then there exists  $i'$  such that two bottom rows of the following diagram is a commutative diagram of cofibrations

$$\begin{array}{ccccccc}
 & & \Sigma^{jq+1} V(0) & \xrightarrow{\lambda'' = \lambda} & \Sigma^{jq+1} W & = & \Sigma^{jq+1} M(p^2) \\
 & & \uparrow \pi & & \uparrow \pi' & & \\
 \Sigma^{-1} V_j & \xrightarrow{\delta} & V_j & \xrightarrow{\lambda'} & C(\delta_{V_j}) & \xrightarrow{\rho'} & V_j \\
 \uparrow i & & \uparrow i & & \uparrow i' & & \uparrow i \\
 \Sigma^{-1} V(0) & \xrightarrow{\delta_{V(0)}} & V(0) & \xrightarrow{\lambda} & M(p^2) & \xrightarrow{\rho} & V(0)
 \end{array}$$

Since  $\rho_*: BP_* M(p^2) \rightarrow BP_* V(0)$ ,  $i_*: BP_* V(0) \rightarrow BP_* V_j$  are all canonical projection, then  $i'_*$ ,  $\rho'_*$  also are. Then from  $\lambda_* = p$ , we get  $\lambda'_* = p$  and the middle row realizes the following short exact sequence (write  $C(\delta_{V_j}) = M(p^2, v_1^j)$ )

$$0 \rightarrow BP_*/(p, v_1^j) \xrightarrow{p} BP_*/(p^2, v_1^j) \rightarrow BP_*/(p, v_1^j) \rightarrow 0$$

Since  $i'$  realizes the canonical projection  $BP_*/(p^2) \rightarrow BP_*/(p^2, v_1^j)$ , then if  $W$  is the cofibre of  $i'$ , then  $BP_* W = BP_*/(p^2)$  and it follows from the uniqueness that  $W = M(p^2)$  and there is  $\lambda''$  which is compatible with  $\lambda$  such that  $\lambda'' \pi = \pi' \lambda'$  or  $\lambda \pi = \pi' \lambda'$ .

If  $j \leq p^{n-1}$  and  $j \equiv 0 \pmod{p}$ , then  $j = kp^s$  ( $s \geq 1$ ) and it follows from corollary 3.13 that  $V_j$  is a split ring spectrum over  $Z_p$ . In the proof of Theorem I, we have had a self map

$$\eta_j: \Sigma^{p^{n-1}(p+1)q} V_j \rightarrow V_j$$

such that  $(\eta_j)_* = v_2^{p^{n-1}}$  and we may assume  $\eta_j \in \mathcal{C}_*(V_j)$  since its components in  $\delta \mathcal{C}_*$ ,  $\bar{h} \mathcal{C}_*$ ,  $\bar{h} \delta \mathcal{C}_*$  will induce zero homomorphism. Then it follows from (3.19) that  $\delta \eta_j^p = \eta_j^p \delta$  and then

there exists  $\eta'$  such that the following diagram commutes

$$\begin{array}{ccccc}
 \Sigma^{-1} V_j & \xrightarrow{\delta} & V_j & \xrightarrow{\lambda'} & M(p^2, v_1^j) = C(\delta_{v_j}) \\
 \uparrow \eta_j^p & & \uparrow \eta_j^p & & \uparrow \eta' \\
 \Sigma^{tp^n(p+1)q} V_j & \xrightarrow{\delta} & \Sigma^{tp^n(p+1)q} V_j & \xrightarrow{\lambda'} & \Sigma^{tp^n(p+1)q} M(p^2, v_1^j)
 \end{array}$$

Then

$$\begin{aligned}
 \beta_{tp^n/j} &= \tau \pi \eta_j^p i j_0 = \tau' \lambda \pi \eta_j^p i j_0 = \tau' \pi' \lambda' \eta_j^p i j_0 = \tau' \pi' \eta' \lambda' i j_0 \\
 &= \tau' \pi' \eta' i' \lambda j_0 = \tau' \pi' \eta' i' j' p,
 \end{aligned}$$

where we use the equation  $\tau = \tau' \lambda$ ,  $\lambda j_0 = j' p$  which are easily seen from the following commutative diagram

$$\begin{array}{ccccc}
 S^o & \xrightarrow{p} & S^o & \xrightarrow{j_0} & V(0) & \xrightarrow{\tau} & \Sigma S^o \\
 \parallel & & \downarrow p & & \downarrow \lambda & \nearrow \tau' & \\
 S^o & \xrightarrow{p^2} & S^o & \xrightarrow{j'} & M(p^2) & & 
 \end{array}$$

So  $\beta_{tp^n/j}$  are divisible by  $p$ . Theorem II is then proved.

Theorem III can be obtained from Theorem I directly.

*Acknowledgements*—I would like to thank the referee for pointing out a gap and an error in our original manuscript, his comments make this revised version here. Also like to thank professor Zhou Xueguang and Sun Yufeng for their kind encouragement during preparation of this paper.

#### REFERENCES

1. J. F. ADAMS: On the groups  $J(X)$  IV. *Topology* **5** (1965), 21–71.
2. J. F. ADAMS: Stable homotopy and generalized homology. Univ. of Chicago. Lect. Notes 1971.
3. D. C. JOHNSON, H. R. MILLER, W. S. WILSON and R. S. ZAHLER: Boundary homomorphisms in the generalized Adams spectral sequence and the nontriviality of infinitely many  $\gamma_i$  in stable homotopy. Conf. on Homotopy theory (Northwestern Univ. Evanston Ill 1974) (Edited by D. H. Davis), p. 47–59. *Math. Notes and Symposia*, Vol 1. Soc. Mat. Mex. Mexico City 1975.
4. P. S. LANDWEBER: Annihilator ideals and primitive elements in complex cobordism. *Illinois J. Math.* **17** (1973), 273–284.
5. A. LIULEVICIUS: Zeroes of the cohomology of the Steenrod algebra. *Proc. Amer. Math. Soc.* **14** (1963), 972–976.
6. H. R. MILLER and W. S. WILSON: On Novikov's  $\text{Ext}^1$  modulo an invariant prime ideal. *Topology* **15** (1976), 131–141.
7. H. R. MILLER, D. C. RAVENEL and W. S. WILSON: Periodic phenomena in the Adams–Novikov spectral sequence. *Ann. Math.* **106** (1977), 469–516.
8. J. MILNOR: Steenrod algebra and its dual. *Ann. Math.* **67** (1958), 150–171.
9. S. OKA: A new family in the stable homotopy groups of spheres. *Hiroshima Math. J.* **5** (1975), 87–114.
10. S. OKA: Realizing some cyclic  $BP_*$  modules and applications to stable homotopy of spheres. *Hiroshima Math. J.* **7** (1977), 427–447.
11. S. OKA: Small ring spectra and  $p$ -rank of the stable homotopy of spheres. *Contemp. Math.* **V. 19** (1983), 267–308.
12. S. OKA: Multiplicative structure of finite ring spectra and stable homotopy of spheres. *Algebraic Topology* (Aarhus 1982) 418–441. Lect. Notes in Math. **V. 1051** Springer-Verlag 1984.
13. D. C. RAVENEL: The nonexistence of odd primary arf invariant elements in stable homotopy. *Math. Proc. Phil. Soc.* **83** (1978), 429–443.
14. D. C. RAVENEL: A novice's guide to the Adams–Novikov spectral sequence. Proc. Evanston Homotopy theory Conf. Lect. Notes in Math. **V. 658** p. 404–475.
15. L. SMITH: On realizing complex cobordism modules. Applications to the stable homotopy of spheres. *Amer. J. Math.* **92** (1970), 793–856.

16. L. SMITH: On realizing complex cobordism modules IV. *Amer. J. Math.* **99** (1977), 418–436.
17. E. THOMAS and R. ZAHLER: Generalized higher order cohomology operations and stable homotopy groups of spheres. *Advance in Math.* **20** (1976), 287–328.
18. H. TODA: On spectra realizing exterior parts of the Steenrod algebra. *Topology* **10** (1971), 53–65.
19. R. S. ZAHLER: Fringe families in the stable homotopy. *Trans A.M.S.* **224** (1976), 243–254.
20. LIN JINKUN: Second periodicity elements in stable homotopy groups of spheres. *KeXue TongBao (Science Bulletin)* **33** (1988) (23), p. 1924–1927.

*Department of Mathematics*  
*Nankai University*  
*Tianjin*  
*People's Republic of China*